

200200050



TO ALL TO WHOM THESE: PRESENTS SHALL COME;

#### Kansas Agricultural Experiment Station

THEFERS, THERE HAS BEEN PRESENTED TO THE

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED. OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS TROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, IN CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN CLING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY TION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CHARIFFED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE WAS ALST THE SECOND TO USE OF THE SECOND THE

WHEAT. COMMON

'Stanton'

In Testimony Phenol. I have hereunto set my hand and caused the seal of the Hunt Intiety Frotestion Office to be affixed at the City of Washington, D.C. this sixteenth day of September, in the year two thousand two.

gem zhe

Commissioner Plant Variety Protection Office Aaricultural Marketina Service ) . . . . . U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

#### Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426). APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse) 1. NAME OF OWNER 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME 3. VARIETY NAME Kansas Agricultural Experiment Station KS95H167-3 Stanton 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 5. TELEPHONE (Include area code) FOR OFFICIAL USE ONLY PVPO NUMBER Waters Hall 785-532-6147 Kansas State University 00200050 Manhattan KS 66506 6. FAX (include area code) 785-532-6563 FILING DATE IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) 8. IF INCORPORATED, GIVE STATE OF INCORPORATION 9. DATE OF INCORPORATION University 10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) FILING AND EXAMINATION FEES: T. Joe Martin Kansas State University Agricultural Research Center-Hays 1232 240th Avenue Hays, KS 67601-9228 11. TELEPHONE (Include area code) 12. FAX (Include area code) 13. E-MAIL 14. CROP KIND (Common Name) 785-625-3425 785-623-4369 jmartin@oznet.ksu.edu wheat 17. IS THE VARIETY A FIRST GENERATION HYBRID? 15. GENUS AND SPECIES NAME OF CROP 16. FAMILY NAME (Botenical) Triticum aestivum XIXNO Gramineae ☐ YES CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse) DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 83(a) of the Plant Variety Protection Act) a. K Exhibit A. Origin and Breeding History of the Variety XX YES (If "yes", answer items 20 and 21 below) NO (If "no", go to item 22) b. KK Exhibit B. Statement of Distinctness c. KK Exhibit C. Objective Description of Variety 20. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF CLASSES? XIX YES NO d. KK Exhibit D. Additional Description of the Variety (Optional) IF YES, WHICH CLASSES? XX FOUNDATION XXREGISTERED XX CERTIFIED e. KK Exhibit E. Statement of the Basis of the Owner's Ownership f. Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public 21. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? XX YES ☐ NO g. KK Filing and Examination Fee (\$2,705), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office) IF YES, SPECIFY THE FOUNDATION REGISTERED CERTIFIED NUMBER 1.2.3. etc. 11-23-01 (If additional explanation is necessary, please use the space indicated on the reverse.) HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. S, OR OTHER COUNTRIES? 23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? XX YES ☐ NO YES KK NO IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space Indicated on reverse.) IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.) 24. The owners declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the cartificate. The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties. SIGNATURE OF OWNER SIGNATURE OF OWNER NAME (Please print or type) NAME (Please print or type) Dr. Forrest Chumley CAPACITY OR TITLE CAPACITY OR TITLE DATE Associate Director

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository: (4) check drawn on a U.S. bank for \$2,705 (\$320 filing fee and \$2,385 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
  - (1) identify these varieties and state all differences objectively;
  - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
  - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Fall, 2001 sold seed in U.S.

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed/ls-sd.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is astimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gethering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audictape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5984 (voice and TDD). USDA is an equal opportunity provider and employer.

S&T-470 (04-01) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (02-99) which is obsolete.

Stanton, PVP Application

Exhibit A: Origin and Breeding History of the Variety. (Revised)

Stanton was selected from the cross PI220350/KS87H57//TAM200/KS87H66/3/KS87H325

#### Parents:

PI220350 = A Russian wheat aphid resistant accession from Afghanistan.

KS87H57 = PI191545/5\*Larned/Eagle//Sage/3/TAM 105

**TAM 200** 

KS87H66 = Selection made from a randomly mated population. The following  $F_1$ s or lines were randomly mated to produce the population.

- 1. Hawk/4/PI194358/2\*Larned/Eagle//Sage/3/Larned/Trison (F<sub>1</sub> from last cross)
- 2. Tut/4/PI195713/2\*Larned/Eagle//Sage/3/Larned/Trison (F<sub>1</sub> from last cross)
- 3. TAM 105/3/PI195713/2\*Larned/Eagle//Sage (F<sub>1</sub> from last cross)
- 4. TAM 105/3/PI194358/2\*Larned/Eagle//Sage (F<sub>1</sub> from last cross)
- 5. TAM 105/4/ PI194358/2\*Larned/Eagle//Sage/3/Larned/Trison (F<sub>1</sub> from last cross)
- 6. TAM 105/4/PI194358/3/Larned/ Eagle//Sage/4/PI195713/2\*Larned /Eagle//Sage (F<sub>1</sub> from last cross)
- 7. TAM 105/Wings (F<sub>1</sub> from last cross)
- 8. Arkan/Wings (F<sub>1</sub> from last cross)
- 9. Arkan/4/PI195713/2\*Larned/Eagle//Sage/3/Larned/Trison (F<sub>1</sub> from last cross)
- 10 Arkan/4/PI194358/2\*Larned/Eagle//Sage/3/Larned/Trison (F<sub>1</sub> from last cross)
- 11 PI194358/2\*Larned/Eagle//Sage/3/Arkan (F<sub>1</sub> from last cross)
- 12 PI194358/2\*Larned/Eagle//Sage/3/TAM 105//Payne/Amigo (F<sub>1</sub> from last cross)
- 13 PI194358/2\*Larned/Eagle//Sage/3/ND7637 (F<sub>1</sub> from last cross)
- 14 PI194358/2\*Larned/Eagle//Sage/3/ND7637/TX73V169 (TAM 101/Centurk) (F<sub>1</sub> from last cross)
- 15 PI194358/2\*Larned/Eagle//Sage/3/ND7637/TX73V169 (F<sub>1</sub> from last cross)
- 16 PI194358/2\*Larned/Eagle//Sage/3/TAM 105//Larned/Trison (F<sub>1</sub> from last cross)
- 17 PI195713/2\*Larned/Eagle//Sage/3/Larned/Trison (KS80HA845)
- 18 Newton sib (KS75210)

#### KS87H325 = RL6005/RL6008//2\*Larned/Cheney/Larned/4/Bennett sib/5/TAM 107

1990:  $F_1$  was grown in field at Hays, KS and harvested in bulk (all F1 plants harvested together). Segregation was noted for height, maturity, and resistance to leaf rust among the  $F_1$  plants. The only selection criteria used was for resistance to leaf rust.

1991: F<sub>2</sub> grown in field at Hays, KS and harvested in bulk. The F<sub>2</sub> population was observed to segregate for maturity, height, and leaf rust resistance.

1992:  $F_3$  was grown in field at Hays, KS and individual head selections were made. The  $F_3$  population was observed to segregate for the same characteristics as in the  $F_2$ .

#### Exhibit A: Continued (Stanton PVPA)

1993: F<sub>4</sub> head rows were grown at Hays, KS and harvested a single head row after selecting six head selections from the row. Selection criteria used included visual yield estimates, height, maturity, shattering resistance, leaf rust resistance, test weight, and grain hardness.

1994: F<sub>5</sub>, seed from the F<sub>4</sub> head row tested in a preliminary yield test at 3 KS locations. The six reselection head rows were grown at Hays, KS and a single row was harvested and six head selections were made from that single row. Selection criteria used included grain yield and test weight, height, maturity. coleoptile length, bread quality factors as measured with the mixograph, and reaction to Russian wheat aphid, Hessian fly, leaf rust, and wheat streak mosaic virus. Among F<sub>5</sub> selections from this cross the population was segregating for height, maturity ,resistance to Russian wheat aphid, Hessian fly, leaf rust, wheat streak mosaic virus and mixing strength.

1995: F<sub>6</sub>, seed from the F<sub>5</sub> head row was tested in the preliminary yield test at 3 KS locations. The F<sub>6</sub> head selections were grown at Hays and a single row was harvested. Six head selections were harvested from the selected head row. Selection criteria used included grain yield and test weight, height uniformity, uniform maturity, reaction to Russian wheat aphid, Hessian fly, leaf rust, and bread quality factors as measured with the mixograph. Segregation was observed within the line for reaction to Hessian fly and leaf rust. The reselections harvested from the line were homozygous for leaf rust resistance.

1996: F<sub>7</sub> was tested in the Advanced Yield Nursery at 6 KS locations. The F<sub>7</sub> head rows were grown at Hays, KS and a single row was harvested. A small increase plot was grown at Hays using the seed from the selected F<sub>6</sub> head row. Selection criteria included grain Stanton, PVP Application

yield, test weight, kernel size, height, maturity, winter survival, reaction to Hessian fly, stem rust, wheat streak mosaic virus, and hard wheat milling and bread baking tests. We noted no characteristics for which the line was segregating.

1997: F<sub>8</sub> was tested in the Kansas Intrastate Nursery (KIN) at 16 KS locations. A small increase was grown at Hays from the seed harvested from the F<sub>7</sub> head row. 200 head selections were harvested from the increase plot. Selection criteria used included grain yield and test weight, kernel size, height, maturity. shattering, reaction to Russian wheat aphid, Hessian fly, leaf rust, and hard wheat milling and bread baking tests.

1998: F<sub>9</sub> was tested in the KIN, Southern Regional Performance Nursery (SRPN) and the Kansas Performance Test with Winter Wheat Varieties. 200 head rows were grown at Hays and evaluated for uniformity in height, maturity, leaf rust resistance, and resistance to Russian wheat aphid. Selection criteria used on the KIN entries were yield, test weight, kernel size, plant height, maturity, shattering tolerance, reaction to Hessian fly, wheat streak mosaic virus and leaf rust. Plus hard wheat milling and bread baking characteristics.

#### Exhibit A Continued (Stanton PVPA)

999: F<sub>10</sub> tested in same nurseries as in 1998. Breeders seed was produced from the bulked head row seed at Hays, KS. Segregation within the line was not noted. The selection criteria used were the same as in 1998.

2000: F<sub>11</sub> was tested in the Kansas Performance Test with Winter Wheat Varieties. Breeders seed was increased at the Agricultural Research and Extension Center at Colby KS. We noted no segregation within the line. Primary selection criteria were yield, test weight, hard wheat milling and bread baking quality, resistance to leaf rust, wheat streak mosaic virus, Hessian fly, and Russian wheat aphid.

 $2001 \, \mathrm{F}_{12}$  was tested in the same nurseries as in 2000. The breeders seed was planted at Colby KS for production of foundation seed for distribution in the fall of 2001. Segregation within the line was not noted.

Stanton is uniform. Variants are limited to: plants with red glume color that occur at a frequency of less that 1 in 1000 plants and slightly taller plants that occur at a frequency of less than 1 in 1000 plants. The variants as well as the typical plants are commercially acceptable.

Stanton is stable. When sexually reproduced, the variety remains unchanged in its essential and distinctive characteristics. Stanton was observed to be uniform and stable during the four generations prior to release.

Stanton, PVP Application Exhibit B: Statement of Distinctness

Stanton is most similar to TAM 107.

TAM 107 has brown glume color while Stanton has white glumes.

TAM 107 is susceptible to Russian wheat aphid while Stanton is resistant (Appendix A, Table 1).

Leaves of Stanton are distinctly pubescent on the adaxial (upper) surface. Its flag leaf trichomes range from .3 to .5 mm in length. Trichomes on the flag leaves of TAM 107 range from .05 to .075 mm in length (see Appendix A, Figure 1).

Appendix A

Table 1. Relative ratings for resistance to the Russian wheat aphid (RWA) of Stanton and other hard winter wheat cultivars.

	RWA* rating			
Cultivar	1999	2000		
Stanton (KS95H167-3)	4.0	3.3		
Halt	2.3	2.5		
Yumar	3.0	3.3		
Prowers	3.3	3.3		
Prairiered	4.0	3.0		
TAM 107	7.5	7.3		
Jagger	8.0	7.8		
Trego	8.0	8.0		
2137	8.0	7.8		
C.V. (%)	9.6	10.0		
L.S.D. (.05)	0.75	0.59		
		•		

<sup>\*</sup> Entries were rated on a 1-9 scale (1=most resistant). The seedling test was conducted as described by Webser et al. 1987 (J. Econ. Ent. 8:944-949). A randomized complete block design with 4 replications each year was used.

#### Appendix A




Figure 1 - Flag leaves of Stanton and TAM 107 at 45 X.

Public reporting burden for this collection of information is estimated to average 30 minutes reporting the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, ORM, AG Box 7630, Jamie L. Whitten Building, Washington; D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braile, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal opportunity employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE **BELTSVILLE, MD 20705** 

**EXHIBIT C** (Wheat)

9

#### **OBJECTIVE DESCRIPTION OF VARIETY** WHEAT (Triticum spp.)

					Control of the Contro			
NAME OF APPLICANT(S)				FOR OFFICIAL USE ONLY				
Kansas State Univ	versity			PVPO NUMBER				
ADDRESS (Street and No. or RD No., C	Sty, State, and Zip Code)			2002	00050			
Waters Hall				VARIETY NAME				
Kansas State Univ	versity			Stanton				
Manhattan, KS 665				TEMPORARY OR EXPERIME	NTAL DESIGNATION			
				KS95H167-3	1.1			
Place a zero in the first box (e.g a minimum of 100 plants. Cor may be used to determine plant	UCTIONS CAREFULLY: Place 19 19 19 19 19 19 19 19 19 19 19 19 19	number is either 99 or less or sined from varieties entered in	9 or less respectively. I the same trial. Royal H	Data for quantitative plant c	haracters should be based on			
1. KIND:	· · · · · · · · · · · · · · · · · · ·							
	1=Common	2=Durum	3=Club	4=Other	(SPECIFY):			
البيشية			• •					
2. VERNALIZATION:		•		•	-			
2	1=Spring	2=Winter	3=Other (SI	PECIFY):				
3. COLEOPTILE ANT	HOCYANIN:				•			
1	1=Absent	2=Present	•		*			
4. JUVENILE PLANT	GROWTH:							
	1=Prostrate	2=Semi-erect	3=Erect					
5. PLANT COLOR (be	oot stage):	:						
2	1 = Yellow-Green	2 = Green	3 = Blue-Gre	een				
6. FLAG LEAF (boot:	stage):							
1	1 = Erect	2 = Recurved		1 = Not Twisted	2 = Twisted			
7. EAR EMERGENCE								
	Number of Days Earlie	r Than			*			
<u> </u>	·							
0 3	Number of Days Later	Than TAM 107			Page 1 of			

8.	ANTHER COLOR:	
	1 = Yellow 2 = Purple	
9.	PLANT HEIGHT (from soil to top of head, excluding awns)	:
	0 6 cm Taller Than TAM 107	•
	cm Shorter Than	
		* Relative to a PVPO-Approved Commercial Variety Grown in the Same Trial
10.	STEM:	
	A. ANTHOCYANIN	D. INTERNODE (SPECIFY NUMBER)
٠.	1= Absent 2=Present	1 Hollow 2=Semi-solid 3=Solid
	B. WAXY BLOOM	E. PEDUNCLE
	2 1=Absent 2=Present	2 1=Absent 2=Present
	C. HAIRINESS (last internode of rachis)	25 cm Length
	2 1=Absent 2=Present	
11.	HEAD (at Maturity):	
	A. DENSITY	C. CURVATURE
	2=Middense 3= Dense	2 Inclined 3 = Recurved
	B. SHAPE	D. AWNEDNESS
	1 = Tapering 2= Strap 3 = Clavate 4 = Other (SPECIFY):	1 = Awnless 2 = Apically Awnletted 3 = Awnletted 4 = Awned
2.	GLUMES (at Maturity):	
	A. COLOR	C. BEAK
	$\boxed{1}  1 = \text{White} \qquad 2 = \text{Tan}$	1 = Obtuse 2 = Acute
•	3 = Other (SPECIFY):	3 =Acuminate
	B. SHOULDER	D. LENGTH
	3 = Rounded 4 = Square 5 = Elevated 6 = Apiculate	1 = Short 2 = Medium (ca. 7mm) (ca. 8mm) 3 = Long (ca. 9mm)

C. GLUMES (at Maturity) Continued:  E. WIDTH	200200050
1 = Narrow (ca. 3mm) 2 = Medium (ca. 3.5mm) 3 = Wide (ca. 4mm)	m)
SEED:	
A. SHAPE	C. BRUSH
3 1 = Ovate 2 = Oval 3 = Elliptical	2 1=Short 2=Medium 3=Long
	1 = Not Collared 2 = Collared
B. CHEEK	D. CREASE
2 1=Rounded 2=Angular	2 = Width 60% or less of Kernel 2 = Width 80% or less of Kernel 3 = Width Nearly as Wide as Kernel
	2 1 = Depth 20% or less of Kernel 2 = Depth 35% or less of Kernel 3 = Depth 50% or less of Kernel
E. Color	G. PHENOL REACTION (see instructions):
1=White 2= Amber 3= Red 4= OTHER (Specify)	1 = Ivory 2 = Fawn 3 = Light Brown 4 = Dark Brown 5 = Black
F. TEXTURE	
1=Hard 2=Soft	
DISEASE: (0=Not Tested; 1=Susceptible; 2=Resist	ant; 3=Intermediate; 4=Tolerant)
PLEASE INDICATE THE SPECIA	
Stem Rust (Puccinia graminis f. sp. tritici) TRTT RTRQ TPMK QKCS RHMS RTHJ RTQQ	2 Leaf Rust (Puccinia recondita f. sp. tritici) TDBM TCLH PLLM TLLC MGBM PNMR KDBM PLMR
Stripe Rust (Puccinia striiformis)	1 Loose Smut (Ustilago tritici)
Tan Spot (Pyrenophora tritici-repentis)	Flag Smut (Urocystis agropyri)
Halo Spot (Selenophoma donacis)	Common Bunt (Tilletia tritici or T. laevis)
Septoria nodorum (Glume Blotch)	Dwarf Bunt (Tilletia controversa)
Septoria avenae (Speckled Leaf Disease)	Karnal Bunt (Tilletia indica)
Septoria tritici (Speckled Leaf Blotch)	Powdery Mildew (Erysiphe graminis f. sp. tritici)
Scab (Fusarium spp.)	0 "Snow Molds"

14	•					•	N. S. C.	20(	200	050_
14.	Dise	ase (Continued)	(0=Not Tested;	1=Susceptible	; 2=	Resistant;	3=Interm		4=Tolera	
		•	PLEASE IN	DICATE THE	SPECI	FIC RACE	OR STRAI	n testi	ED	•
	1	"Black Point"	(Kernel Smudge)		0	Common l Bipolaris s	Root Rot <i>(</i> pp. <i>)</i>	Fusariun	ı, Cochliob	olus and
	1	Barley Yellow	Dwarf Virus (BYD)	<b>)</b>	0	Rhizocton	ia Root Ro	(Rhizoc	tonia solan	<i>ii)</i>
	1	Soilborne Mos	aic Virus (SBMV)		0	Black Cha	ff <i>(Xantho</i>	monas ca	<i>mpestris</i> pv	v. translucens)
	3	Wheat Yellow	(Spindle Streak) Mos	aic Virus	2	Bacterial I syringae)	eaf Blight	(Pseudoi	monas syrii	rgae pv.
	4	Wheat Streak I	Mosaic Virus (WSM	v) [		Other (SP	ECIFY)			
٠		Other (SPECII	(Y)			Other (SP)	ECIFY)			e e <del>ne</del> rv
		Other (SPECII	FY)	[		Other (SP)	ECIFY)			
		Other (SPECIF	<b>'Y</b> )			Other (SP)	ECIFY)			
15. II	NSECT:	(0=Not Test	ed; 1=Susceptible	2=Resistant	; 3=	-Intermedia	te; 4=To	lerant)		
			PLEASE S	PECIFY BIOT	YPE (v	where neede	d)		٠	
	3		ny <i>etiola destructor)</i> s Biotype (H18)			Other (SPE	CIFY)			·
•	0	Stem Sawfly (Co	ephus spp.)			Other (SPE	CIFY)			
	0	Cereal Leaf Beet	le <i>(Oulema melanop</i>	a) [		Other (SPE	CIFY)			
	2	Russian Aphid	Diuraphis noxia	ſ	7 '	Other (SPE	CIFY)			

Other (SPECIFY)

Other (SPECIFY)

6. ADDITIONAL INFORMATION ON ANY ITEM ABOVE, OR GENERAL COMMENTS

Greenbug (Schizaphis graminum)

Aphids

Stanton, PVP Application Exhibit D: Additional description of the variety


The hard wheat milling properties and bread baking characteristics of Stanton were compared to currently grown varieties by the members of the Wheat Quality Counsil in 1997 and 1998. They rated Stanton's overall baking quality as above average in both years. I have attached copies of their reports ( see Appendix B).

200200050

# Milling and Baking Test Results for

#### Hard Winter Wheats

Harvested in 1997



#### 48th Report on Wheat Ouality

Ascoordinated efforeby the agricultural and baking industries to improve wheat






#### 1997 Miag Multomat (Small-Scale) Samples Kansas

			Stanton	丢			
Sample Number	97-407	97-408	97-409	97-410	97-419	97-420	
Variety Identification	Lamed	KS95	KS95	KS96	Jagger	KS84	
	(check)	H176-1	H167-3	HW62-6	(check)	063-2W	
Wheat Data							
FGIS Classification	HRW	HRW	HRW	HWW 2.4%HR	HRW	HWW	
U.S. Bushel Weight (lbs)	60.2	59.0	59.7		60.1	60.9	
Hectoliter Weight (kg)	77.5	75.9	76.8	79.5	77.3	78.4	
1000 Kernel Weight	36.0	29.2	37.4	33.6	23.0	26.5	
NIR Hardness	67	65	67	74	71	72	
Wheat Size Test			_		-		
Over 7 Wire (%)	82.8	68.2	88.3		51.6	61.7	
Over 9 Wire (%)	17.1	31.7	11.7		47.6	38.1	j j
Through 9 Wire (%)	0.0	0.2	0.1	0.1	0.8	0.2	
Single Kernel							
Hardness	59	59	58	1		76	
Weight (mg)	34.9	29.4	37.7	33.7	29.4	29.6	
Diameter (mm)	2.65	2.47	2.88	2.60	2.53	2.49	Ī
Moisture (%)	10.80	10.50	10.40	10.20	9.60	9.80	
Protein (% *)	11.60	12.30	12.20	12.20	14.30	13.90	
Ash (% *)	1.34	1.34	1.19	1,29	1.22	1.24	<u> </u>
Milling and Straight	t Grade Flo	ur Data					
Straight Grade					70.0	60.4	
Flour Yield (% T.P.)	73.0	73.8	73.0	72.0	72.3	69.4	
Moisture (%)	13.70	13.60	13.60	13.50	13.20	13.20	
Protein (% *)	10.60	11.00	11.20		13.30	13.30	
Ash (% *)	0.43	0.40	0.38	0.42	0.43	0.42	
Glutomatic	<del>                                     </del>						
Wet (% *)	2.85	2.75	2.76	2.93	3.59	3.51	
Dry (% *)	0.96	0.98	0.98	1.01	1.23	1.21	
Index	95.4	98.7	98.6	98.5	98.2	98.3	
Solor							
Kent Jones/Simon	-0.38	0.79	-0.46	-0.28	-0.40	0.79	
alling Number *	445	506	402	405	592	561	
Starch Damage	<del>  </del>						
Verage Micron Size	+				•		
Fisher S S	10 6	10 2	10 6	10.7	10 3	10 6	


## 1997 Physical Testing Results (Small Scale) Kansas






Abs: 62.5 %, Peak: 14.5 min. Stab: 29.75 min.

97-419 Jagger (check)




#### SPONGE CHARACTERISTICS

(Small Scale) Kansas




#### **BAKE ABSORPTION**


(Small Scale) Kansas




# BAKE MIX TIME (Small Scale) Kansas



#### MIXING TOLERANCE



# DOUGH CHAR. 'OUT OF MIXER' (Small Scale) Kansas




# DOUGH CHAR. 'OUT OF MIXER', DESCRIBED (Small Scale) Kansas

	Sticky	Wet	Tough	Good	Excellent
97-407 Larned (check)	2	0	0	12	1
97-408 KS95H176-1	2	0	2	11	0
97-409 KS95H167-3	0	0	1	12	2
97-410 KS95HW62-6	1	. 1	0	12	1
97-419 Jagger (check)	1	0	5	9	0
97-420 KS84063-2W	1	0	3	11	0

Frequency Table

# DOUGH CHAR. 'AT MAKE UP' (Small Scale) Kansas




# DOUGH CHAR. 'AT MAKE UP', DESCRIBED (Small Scale) Kansas

	Sticky	Wet	Tough	Good	Excellent
97-407 Larned (check)	1	0	0	13	1
97-408 KS95H176-1	0	0	2	13	0
97-409 KS95H167-3	0	0	1	14	0
97-410 KS95HW62-6	0	0	1	13	1
97-419 Jagger (check)	0	0	7	8	0
97-420 KS84063-2W	0	0	5	9	1

Frequency Table

#### CRUMB GRAIN (Small Scale) Kansas




# CRUMB GRAIN, DESCRIBED (Small Scale) Kansas

	Open	Dense	irregular
97-407 Larned (check)	6	6 1	
97-408 KS95H176-1	97-408 95H176-1 4		5
97-409 KS95H167-3	5	2	4
97-410 KS95HW62-6	3	2	6
97-419 Jagger (check)	4	4	3
97-420 KS84063-2W	3	3	6

Frequency Table

200200050

# CRUMB TEXTURE (Small Scale) Kansas




# CRUMB TEXTURE, DESCRIBED (Small Scale) Kansas

	Coarse	Harsh	Silky
97-407 Larned (check)	3	3	5
97-408 KS95H176-1	4	4	3
97-409 KS95H167-3	4	2	5
97-410 KS95HW62-6	3	3	5
97-419 Jagger (check)	3	3	6
97-420 KS84063-2W	97-420 34063-2W 4 0		8

Frequency Table

# CRUMB COLOR (Small Scale) Kansas




#### CRUMB COLOR, DESCRIBED

(Small Scale) Kansas


	Yellow	Gray	Dull	Сгевту	Bright White
97-407 Larned (check)	2	0	0	11	1
97-408 KS95H176-1	1	0	1	11	1
97-409 KS95H167-3	1	0	0	13	0
97-410 KS95HW62-6	1	0	1	10	2
97-419 Jagger (check)	2	0	2	8	2
97-420 KS84063-2W	0	0	1	7	6

Frequency Table

## LOAF VOLUME (Small Scale) Kansas



#### OVERALL BAKING QUALITY



#### COOPERATOR'S COMMENTS

(Small Scale) Kansas

#### 97-407 Larned (check)

- •SL CR BRIGHT
- 4' had sl higher vol.but was sticky, more open grain
- •MARGINAL STRENGTH, AVERAGE VOLUME
- · Slightly open, slightly irregular cells. Dry-firm body.
- good volume
- •63 ppm AA; uniform crumb grain; rough crust.
- •GOOD MIXING TOLERANCE.

  MOISTURE 13.68 ASH .436 (14%MB) PROTEIN 10.43 (14%MB)
  FARINOGRAPH ABS 62.8 (14%MB) PEAK 4.0 MTI 20
  FALLING NUMBER 393 STARCH DAMAGE 5.94
- Very good loaf volume for its flour protein content.

#### 97-408 KS95H176-1

- •SL CR BRIGHT VERY OPEN GRAIN
- dough was better at 9' but had lower vol.
- •VERY STRONG MIXING DOUGH, OPEN GRAIN, GOOD VOLUME
- Slightly irregular cells. Dry, very firm body Dry, dead doughs at make-up
- · low absorption
- •13 ppm AA; fine uniform cells; slightly rough crust.
- •LOW ABSORPTION.

  MOISTURE 13.55 ASH .423 (14%MB) PROTEIN 10.82 (14%MB)

  FARINOGRAPH ABS 56.0 (14%MB) PEAK 2.5 MTI 15

  FALLING NUMBER 411 STARCH DAMAGE 7.89
- Very good loaf volume for its flour protein content.
   Good crumb and texture.

#### 97-409 KS95H167-3

- at 7.5' had very shrunken sides; at 9' was more open
- GOOD STRENGTH, EXCELLENT VOLUME, SL. OPEN GRAIN
- · Creamy, slightly irregular cells
- •13 ppm AA; thin cell walls; exc. crumb grain.
- •LOW ABSORPTION.

  MOISTURE 13.64 ASH .423 (14%MB) PROTEIN 10.85 (14%MB)

  FARINOGRAPH ABS 55.6 (14%MB) PEAK 8.5 MTI 10

  FALLING NUMBER 354 STARCH DAMAGE 6.78
- Very good loaf volume for its flour protein content.
   Good crumb and texture.

#### COOPERATOR'S COMMENTS--Cont'd

(Small Scale) Kansas

#### 97-410 KS95HW62-6

- •best vol.@5'; sl better grain @4'; sl sticky @ 6'
- •GOOD VOLUME, NICE INTERIOR
- · Creamy, slightly irregular cells
- •38 ppm AA;
- •GOOD MIXING TOLERANCE.

  MOISTURE 13.55 ASH .441 (14%MB) PROTEIN 10.87 (14%MB)
  FARINOGRAPH ABS 58.9 (14%MB) PEAK 6.0 MTI 10
  FALLING NUMBER 378 STARCH DAMAGE 8.14
- · Good loaf volume for its flour protein content.

#### 97-419 Jagger (check)

- VERY OPEN SHOTTY GRAIN
- •7' & 10' mix were more elastic and had lower vol.
- Strong/Slightly Creamy
- •VERY GOOD STRENGTH, GOOD VOLUME
- · Irregular, slightly open cells. Volume down. Silky texture
- good protein low absorption
- 38 ppm AA;slightly sticky out of mixer.
- •GOOD MIXING TOLERANCE. GOOD MIX TIME.

  MOISTURE 12.98 ASH .456 (14%MB) PROTEIN 12.96 (14%MB)

  FARINOGRAPH ABS 60.8 (14%MB) PEAK 6.5 MTI 10

  FALLING NUMBER 483 STARCH DAMAGE 8.14
- Crumb color was yellow and dull
- Good loaf volume for its flour protein content.
   Good crumb and texture.

#### 97-420 KS84063-2W

- •VERY OPEN SHOTTY GRAIN
- •7' was very shrunken; 11' had low vol & stringy dough
- Overall the Best
- •GOOD CRUMB COLOR, ACCEPTABLE INTERIOR, ABOVE AVG. VOLUME
- Irregular, slightly open cells. Good body. Silky texture
- good color good protein
- •25 ppm AA; exc. loaf symmetry and crumb grain; soft crumb.
- •GOOD FLOUR OVERALL.

  MOISTURE 12.88 ASH .429 (14%MB) PROTEIN 12.83 (14%MB)
  FARINOGRAPH ABS 60.5 (14%MB) PEAK 8.5 MTI 10
  FALLING NUMBER 420 STARCH DAMAGE 7.45

2002000505



# 1998 Milling and Baking Test Results for Hard Winter Wheats

#### Editor:

Patrick J. McCluskey
Department of Grain Science
Kansas State University
Manhattan, KS

#### **Coordinator:**

Ben Handcock, Executive Vice President Wheat Quality Council Pierre, SD

#### Kansas: 1998 (Small-Scale) Samples


·· ·			Stanton					
Sample Number	98-404	98-405	98-406	98-407	98-408	98-409	98-410	98-411
Variety Identification	Karl 92	Lamed	KS95	KS95	KS96	KS96 <sub>5</sub>	KS84	KS85
	(check)	(check)	H167-3	HW62-6	HW94	HW118	063-2W	W663-42W
Wheat Data		. "						1
Tineat Data	•						*.	
FGIS Classification	HRW	HRW	HRW	HDWH	HDWH	HDWH	HDWH	HDWH
			1.1% HDWH		0.4% HRW	0.9% HRW	<u></u>	0.6% HRW
Test Weight (lb/bu)	59.8	60.5	59.6	59.7	60.1	60.4	59.3	60.4
Hectoliter Weight (kg/hl)	78.7	79.6	78.4	78.6	79.1	79.5	78.0	79.5
1000 Kernel Weight (gm)	26.8	28.4	28.4	26.3	24.2	29.5	25.7	25.4
NIR Hardness	57	74	67	74	67	54	72	51
Wheat Size Test								
Over 7 Wire (%)	40.2	52.1	56.8	40.4	32.4	56.2	44.7	34.6
Over 9 Wire (%)	59.2	47.1.	42.9	58.6	66.1	43.5	54.8	64.7
Through 9 Wire (%)	0.6	0.8	0.3	1.0	1.5	0.4	0.5	0.7
Single Kernel Analysis					·			
Hardness	64	76	69	81	81	69	78	60
Weight (mg)	28.1	30.3	29.9	28.2	26.0	30.3	26.2	26.1
Diameter (mm)	2.32	2.44	2.44	2.30	2.28	2.44	2.21	2.25
Moisture (%)	10.2	10.4	10.2	10.4	10.3	10.5	10.4	10.4
Protein (%)**	13.6	13.6	12.9	13.5	13.9	13.1	15.1	14.4
Ash (%)*	1.40	1.48	1.43	1.47	1.49	1.38	1.45	1.33
Miling and Straight	Grade Fl	our Data						
	98-404	98-405	98-406	98-407	98-408	98-409	98-410	98-411
Straight Grade		-						
Flour Viold (%T P )	74.7	74 1	74 n	742	746	75.9	71.6	73.6

	98-404	98-405	98-406	98-407	98-408	98-409	98-410	98-411
Straight Grade		-		-				
Flour Yield (%T.P.)	74.7	74.1	74.0	74.2	74.6	75.9	71.6	73.6
Moisture (%)	13.7	13.5	13.2	14.1	13.8	13.5	14.1	14.0
Protein (%)*	12.4	12.3	11.7	12.1	12.3	12.0	14.2	13.5
Ash (%)*	0.46	0.46	0.44	0.44	0.44	0.41	0.46	0.41
Glutomatic							,	
Wet (%)*	30.2	34.5	30.5	31.9	26.9	31.5	37.0	34.7
Dry (%)*	11.4	12.0	10.8	11.2	10.0	10.7	12.9	12.4
Index	98.5	95.4	97.4	96.4	99.3	98.7	97.7	97.7
Color	,							
Agtron Flour Color	85	82	84	83	83	89	82	89
Simon Flour Color	-0.13	0.45	0.32	-0.19	0.09	-0.54	-0.71	-0.93
Falling Number* (sec)	561	633	437	559	567	532	530	478
Avg. Micron Size							<u> </u>	
Fisher Sub Sieve Sizer	19.5	22.9	21.3	23.7	21.9	20.5	22.5	20.2


<sup>\*\* 12%</sup> moisture basis; \*14% moisture basis

#### **Physical Dough Tests**


#### 1998 (Small Scale) Kansas




Mixograms




Abs: 58.7%, Peak: 25.5, Stab: 40.0




Abs: 63.6%, Peak: 5.4, Stab: 5




Abs: 62.9%, Peak: 6.5, Stab: 11.0







Abs: 64.0%, Peak: 2.6, Stab: 4




Abs: 57.6%, Peak: 12.5, Stab: 24.0






Abs: 63.0%, Peak: 4.5, Stab: 4



98-407 KS95HW62-6




Abs: 64.0%, Peak: 3.9, Stab: 4

#### **Physical Dough Tests** 1998 (Small Scale) Kansas

**Farinograms** 


Mixograms




Abs: 61.8%, Peak: 32.0, Stab: 35.5




Abs: 65.0%, Peak: 7.4, Stab: 5




Abs: 57.4%, Peak: 10.0, Stab: 21.5




Abs: 64.0%, Peak: 3.9, Stab: 3




Abs: 62.1%, Peak: 10.5, Stab: 24.0



Abs: 66.7%, Peak: 4.0, Stab: 3



98-411 KS85W663-42W



Abs: 65.4%, Peak: 4.4, Stab: 3

#### SPONGE CHARACTERISTICS

(Small Scale) Kansas

Variety order by rank sum.


Samples with the same letter not different at 5.0% level of significance.

ncoop= 8 chisq= -3.06 chisqc= 24.92 cvchisq= 14.07 crdiff= 12.99

ncoop= 15

cvchisq= 14.07


chisq= 47.01 chisqc= 66.10



#### **BAKE ABSORPTION**

(Small Scale) Kansas

Variety order by rank sum.



# BAKE ABSORPTION, ACTUAL (14% MB) (Small Scale) Kansas

	Coop.	Coop. B	Coop.	Coop. Coop. C D	Соод П	Coop.	Coop. G	Coop. Coop. Coop. Coop. Coop. Coop. G	Coop.	Coop.	00 ⊼	Coop. L		Coop.	Coop.	Coop.	
98-404 Karl 92 (check)	59.0	62.0	58.6	59.0	66.4	62.9	59.0	59.0	68.3	6.69	58.2	63.6	64.0	59.6	62.5	60.6	
98-405 Larned (check)	63.5	62.0	59.0	62.0	65.2	62.7	61.0	59.0	68.4	66.7	63.8	64.0	63.5	63.5	63.0	61.1	
98-406 KS95HW167-3	59.5	61.0	58.0	59.0	63.8	60.3	60.0	59.0	67.2	65.3	59.9	63.0	62.0	2.73	62.0	59.9	
98-407 KS95HW62-6	62.5	62.0	59.0	60.0	6.99	63.5	62.0	59.0	68.9	65.8	58.6	64.0	64.5	62.5	63.0	61.6	
98-408 KS96HW94	60.0	62.0	60.0	60.0	69.0	66.0	61.0	59.0	69.5	67.2	6.09	65.0	65.0	63.0	64.0	62.4	
98-409 KS96HW115	60.5	62.0	59.0	59.0	65.1	63.6	61.0	59.0	68.3	65.6	58.1	64.0	63.5	58.4	63.0	61.1	
98-410 KS84063-2W (Betty)	63.0	64.0	61.7	60.0	68.8	66.4	64.0	61.0	71.4	69.8	59.6	66.7	67.0	64.7	65.0	63.1	
98-411 KS85W663-42W (Heyne)	61.0	64.0	60.4	59.0	65.7	64.1	60.0	60.0	70.2	66.2	57.0	65.4	63.5	59.5	64.5	61.5	

Raw Data

# BAKE MIX TIME, ACTUAI (Small Scale) Kansas

:	Coop.	C80p.	000 000	Coop. Coop. ( D E	О С С	200p.	Coop. G	С Эр Т	Coop. Coop. Coop.	Coop	0 8 7 9	Coop	000 2002 2002	Coop	Coop. Coop.	. Coop.
98-404 Karl 92 (check)	60.0	9.0	6.0	22.0	9.5	6.3	7.2	20.0	0.6	6.9	13.0	8.20	i 0	8.4	21.0	8,4
98-405 Lamed (check)	33.0	6.0	5.0	19.0	3.8	2.8	4.0	7.0	5.0	3.1	5.0	3.0	5.0	2.5	11.0	2.5
98-406 KS95HW167-3	35.0	9.0	6.0	19.0	6.0	4.2	5.6	9.0	4.0	5.0	8.0	4.3	9.0	3.8	9.0	3.5
98-407 KS95HW62-6	32.0	6.0	7.0	20.0	5.5	3.8	5.2	11.0	4.0	2,4	8.0	4.5	6.0	3.5	10.0	3.5
98-408 KS96HW94	60.0	9.0	7.0	20.0	13.0	8.0	10.2	20.0	11.0	8.7	20.0	7.8	25.0	6.5	25.0	6.8
98-409 KS96HW115	4.0	6.0	6.5	20.0	5.	4.2	5.0	16.0	6.0	4.2	7.0	4.0	7.0	3.3	10.0	3.0
98-410 KS84063-2W (Betty)	60.0	6.0	7.5	20.0	5.5	4.2	5.2	13.0	6.0	4.2	11.0	4.6	12.0	3.8	13.0	3.3
98-411 KS85W663-42W (Heyne)	60.0	9.0	ις 10	20.0	6.0	4.8	5.0	12.0	7.0	4.7	8.0	4.5	16.0	3.3	9.0	3.0
	!				10000		A		Annual Property			_	-	-	-	_


#### **BAKE MIX TIME**

(Small Scale) Kansas

Variety order by rank sum.

Samples with the same letter not different at 5.0% level of significance.






#### MIXING TOLERANCE

(Small Scale) Kansas


Variety order by rank sum.

Samples with the same letter not different at 5.0% level of significance.



ncoop= 14 .

chisq= 39.75



Starten


#### DOUGH CHAR. 'OUT OF MIXER'

(Small Scale) Kansas

ncoop= 15 chisq= 7.34 chisqc= 13.60 cvchisq= 14.07 crdiff=

Variety order by rank sum.

No samples different at 5.0% level of significance.



#### DOUGH CHAR. 'OUT OF MIXER', DESCRIBED

(Small Scale) Kansas

•	Sticky	Wet	Tough	Good	Excellent
98-404 Karl 92 (check)	0	0	6	8	1
98-405 Larned (check)	0	1	3	10	1
98-406 KS95HW167-3	0	1	3	10	1
98-407 KS95HW62-6	3	0	3	9	0
98-408 KS96HW94	1	0	8	5	1
98-409 KS96HW115	1	0	3	9	2
98-410 KS84063-2W (Betty)	1	0	5	7	2
98-411 KS85W663-42W (Heyne)	0	0	4	7	4


Frequency Table

# DOUGH CHAR. 'AT MAKE UP'

(Small Scale) Kansas

Variety order by rank sum.

nccop= 16 chisq= 19.23 chisqc= 25.51 cvchisq= 14.07 crdiff= 21.91



# DOUGH CHAR. 'AT MAKE UP', DESCRIBED

(Small Scale) Kansas

	Sticky	Wet	Tough	Good	Excellent
98-404 Karl 92 (check)	0	0	7	5	3
98-405 Larned (check)	0	0	1	13	0
98-406 KS95HW167-3	0	1	4	8	2
98-407 KS95HW62-6	0	0	2	12	1
98-408 KS96HW94	0	0	12	3	0
98-409 KS96HW115	0	0	3	10	2
98-410 KS84063-2W (Betty)	0	0	6	6	3
98-411 KS85W663-42W (Heyne)	0	0	5	7	3


# **CRUMB GRAIN**

(Small Scale) Kansas

ncoop= 16 chisq= 7.44 chisqc= 9.01 cvchisq= 14.07 crdiff=

Variety order by rank sum.

No samples different at 5.0% level of significance.



# CRUMB GRAIN, DESCRIBED


(Small Scale) Kansas

	Open	Dense	Irregular
98-404 Karl 92 (check)	5	4	3
98-405 Larned (check)	8	3	2
98-406 KS95HW167-3	5	3	4
98-407 KS95HW62-6	7	3	4
98-408 KS96HW94	5	5	2
98-409 KS96HW115	4	4	3
98-410 KS84063-2W (Betty)	8	4	2
98-411 KS85W663-42W (Heyne)	8	5	1

Frequency Table

#### **CRUMB TEXTURE**

(Small Scale) Kansas



2

98-408

98-405

98-409

98-411

98-407

98-404

98-410

Stanton 98-406

KS96HW94

Larned (check)

KS85W663-42W (Heyne)


KS96HW115

KS95HW62-6

KS95HW167-3

Karl 92 (check)

KS84063-2W (Betty)



ncoop= 16

r sum= 65.00 mean= 3.69 r sum= 67.50 mean= 3.63

r sum = 73.50mean= 3.57 r sum= 75.00

mean= 3.78 r sum= 77.00

mean= 4.13 r sum= 95.50

5

**VERYHARSH** SILKY CRUMB TEXTURE, DESCRIBED

Cooperator Means

(Small Scale) Kansas

	Coarse	Harsh	Silky
98-404 Karl 92 (check)	3	2	8
98-405 Larned (check)	2	5	7
98-406 KS95HW167-3	4	3	6
98-407 KS95H <b>W</b> 62-6	1	4	9
98-408 KS96HW94	4	5	5
98-409 KS96H <b>W</b> 115	3	3	7
98-410 KS84063-2W (Betty)	2	3	9
98-411   KS85W663-42W (Heyne)	3	2	8

### **CRUMB COLOR**

(Small Scale) Kansas


Variety order by rank sum.

ncoop= 16 chisq= 40.35

chisqc= 51.26

crdiff= 18.77

cvchisq= 14.07



## CRUMB COLOR, DESCRIBED

(Small Scale) Kansas

	Yellow	Gray	Dull	Creamy	Bright White
98-404 Karl 92 (check)	2	0	1	5	6
98-405 Larned (check)	5	0	1	7	0
98-406 KS95HW167-3	5	0 .	4	5	0
98-407 KS95HW62-6	3	0	3	7	1
98-408 KS96H <b>W</b> 94	4	0	2	6	2
98-409 KS96HW115	2	0	0	9	2
98-410 KS84063-2W (Betty)	1	0	0	6	6
98-411   KS85W663-42W (Heyne)	4	0	1	7	1

# CELL SHAPE, DESCRIBED

(Small Scale) Kansas

	Round	Irregular	Elongated
98-404 Karl 92 (check)	0	9	5
98-405 Larned (check)	2	8	4
98-406 KS95HW167-3	1	10	2
98-407 KS95HW62-6	3	9	1
98-408 KS96HW94	2	10	2
98-409 KS96HW115	3	6	5
98-410 KS84063-2W (Betty)	1	7	6
98-411 KS85W663-42W (Heyne)	1	8	4

Frequency Table

# CELL THICKNESS, DESCRIBED

(Small Scale) Kansas

	Too Thin	Too Thick	Variable	Broken	Acceptable
98-404 Karl 92 (check)	1	0	6	0	6
98-405 Larned (check)	0	3	4	0	6
98-406 KS95HW167-3	0	2	8	0	4
98-407 KS95HW62-6	0	3	6	0	4
98-408 KS96HW94	1	3	4	0	5
98-409 KS96HW115	0	2	6	0	5
98-410 KS84063-2W (Betty)	0	2	7	0	4
98-411   S85W663-42W (Heyne)	0	2	5	0	6

# LOAF WEIGHT, ACTUAL (Small Scale) Kansas

	Coop. C	Соор. В	င်္လ လ	Coop. Coop. C D	Соор. П	Coop. Coop.	იმე	Coop. H	Coop. Coop. Coop. Coop. Coop. Coop. Coop. Coop. Coop.	Coop.	00 7 7	Coop.	Coop.	Coop.	Coop.	
98-404 Karl 92 (check)	466.0		490.0	,	153.4	139.0	132.4	421.0	454.0	150.5	597.0	139.7	460.2	130.0	421.2	131.6
98-405 Larned (check)	456.0		495.0		152.5	139.8	136.3	423.0 448.0	448.0	154.0	594.0	141.2	461.1	135.0	418.5	133.7
98-406 KS96HW167-3	462.0		490.0		149.1	136.7	136.1	420.0	449.0	150.0	594.0	139.7	459.6	129.0	129.0 417.0	133.9
98-407 KS95HW62-6	457.0		490.0		153.0	138.6	136.0	419.3	448.0	150.8	597.0	142.7	461.4	133.0	416.8	137.2
98-408 KS96HW94	464.0		495.0		151.8	139.2	136.3	422.5	453.0	151.2	605.0	144.6	460.8	133.0	422.7	134.8
98-409 KS96HW115	462.0		495.0		152.0	140.4	136.1	420.5 448.0	448.0	153.4	598.0	145.6	464.3	132.0	419.1	137.4
98-410 KS84063-2W (Betty)	457.0		495.0		152.9	139.2	142.1	422.0	451.0	154.6	596.0	144.7	459.8	135.0	417.8	138.5
98-411 85W663-42W (Heyne)	460.0		495.0		151.0	138.2	136.6	420.0	446.0	149.4 594.0	594.0	146.2	463.0	132.0	132.0 418.9	136.2

# LOAF VOLUME, ACTUAI (Small Scale) Kansas

		÷.						
Coop.	900	800	855	795	840	825	865	890
Coop.	2226	2133	2256	2250	2103	2256	2183	2153
Coop.	995	915	950	925	965	925	1060	1025
Coop.	3000	2800	2875	2725	2500	2725	2913	2825
Coop.	1025	925	920	875	950	945	1100	975
Coop.	2625	2350	2450	2375	2600	2300	2425	2450
Coop.	955	1020	1010	925	915	995	1045	1025
Coop.	2625	2550	2500	2325	2275	2425	2375	2550
Соор Н	2800	2800	2800	2800	2700	2700	2800	2750
Соор. G	970	940	930	915	955	935	066	1020
Соор.	955	006	965	915	915	096	1105	1000
Соор. Е	970	006	975	970	950	925	1005	975
Coop.	3045	2809	2839	2853	2750	2750	2927	2868
လ လ ပ	2850	3125	3125	3050	2475	2975	3300	2900
Coop. C	1125	066	1095	1045	900	1045	980	1055
Coop.	2600	2600	2625	2600	2550	2650	2675	2775
	98-404 Karl 92 (check)	98-405 Larned (check)	98-406 KS95HW167-3	98-407 KS95HW62-6	98-408 KS96HW94	98-409 KS96HW115	98-410 KS84063-2W (Betty)	98-411 KS85W663-42W (Heyne)


#### LOAF VOLUME

(Small Scale) Kansas

Variety order by rank sum.

Samples with the same letter not different at 5.0% level of significance.

ncoop= 15 chisq= 20.06 chisqc= 25.45 cvchisq= 14.07 crdiff= 21.54





Cooperator Means

(Small Scale) Kansas

Variety order by rank sum.

KS96HW94

KS95HW62-6

Larned (check)

KS84063-2W (Betty)

KS85W663-42W (Heyne)

KS96HW115

Karl 92 (check)

98-406 bc KS95HW167-3

**VERY POOR** 

. 98-408

98-407

98-405


98-409

98-410 c

98-404 G

Starton

Samples with the same letter not different at 5.0% level of significance.



ncoop= 15 chisg= 15.39 chisqc= 18.16

5

cvchisq= 14.07 crdiff= 23.34

r sum= 54.00 r sum= 63.50

#### COOPERATOR'S COMMENTS

#### (Small Scale) Kansas

#### 98-404 Karl 92 (check)

- Very strong dough
- crumb grain=sl. dense; cell shape = sl. elongated; color=white
- strong mixing flour, irregular streaky grain; good volumes, creamy crumb color
- Long mix time
- 13 ppm AA
- Slightly weak crumb texture
- MOISTURE 13.43 ASH(14%) 0.489 PROTEIN(14%) 12.21
- Bake mix time may be too long
- Best of show in quality, but low absorption; cell thickness=thin=desirable
- Slight Shell Top, Break Little Rough, Slightly Dry
- v. long mixt; hi abs.; v. nice grain; elastic dough; v. hi volume; crumb texture=good; crumb grain=closed
- Would be a good flour to blend with weaker flours.

#### 98-405 Larned

- Dead dough
- crumb grain=sl. dense; cell shape = sl. elongated
- strong flour, open, irregular grain, low volumes, creamy crumb color
- Good absorption and crumb grain
- 75 ppm AA; oxidation; mix time; volume
- Slightly weak crumb texture
- MOISTURE 13.29 ASH(14%) 0.461 PROTEIN(14%) 12.07
- sl. short mixt; avg. abs.; poor grain; yellow; weak dough; good volume; at makeup=weak

#### 98-406 KS95H167-3

- · dead dough, lower absorption, good internals
- crumb grain=si. dense; cell shape = si. elongated
- good mixing strength, open grain, slightly low to average volumes.
- Good absorption, loaf vol. and crumb grain
- 38 ppm AA
- Strong and extensible dough out of the mixer
- MOISTURE 13.02 ASH(14%) 0.483 PROTEIN(14%) 11.64
- Very good loaf volume for its flour protein content. Very good crumb grain.
- out of mixer=sl. tough; cell thickness=thin=desirable
- Nice Break, Little Dry
- avg. mixt; avg. abs.; sl. open grain; yellow; elastic dough; good volume.

# COOPERATOR'S COMMENTS (con't.)

#### (Small Scale) Kansas

#### 98-407 KS95HW62-6

- crumb grain=sl. dense; cell shape = sl. elongated; color=white
- strong mixing flour, open grain, average volume
- Good absorption, loaf vol. and crumb grain
- 38 ppm AA
- Slightly weak crumb texture
- MOISTURE 13.85 ASH(14%) 0.458 PROTEIN(14%) 11.97
- out of mixer = very tough
- Break Slightly Rough, Little Dry
- avg. mixt; hi abs.; avg. grain; silky texture; nice dough; avg. volume.

#### 98-408 KS96HW94

- slight core, very strong dough, good internals
- tough, bucky
- out of mixer=tough/old
- very strong mixing flour, open irregular grain, low volume
- Long mix time- messy dough while mixing- otherwise good
- 0 ppm AA; mix time
- MOISTURE 13.42 ASH(14%) 0.430 PROTEIN(14%) 12.13
- Crumb grain somewhat questionable, mix time too long.
- Too long of mix times; cell thickness=thin=desirable
- Good Break, Dry; Long time for mix pickup added .5 min
- v. long mixt, hi abs.; nice grain; tough sponge; elastic dough; low volume.
- Sides of loaf were a little concave.

#### 98-409 KS96HW115

- slightly dead dough
- crumb grain=sl. dense; cell shape = sl. elongated; color=sl. creamy
- mix strength was very good, open, irregular grain, low volumes
- Good absorption, loaf vol. and crumb grain
- 38 ppm AA
- Slightly weak crumb texture
- MOISTURE 13.15 ASH(14%) 0.419 PROTEIN(14%) 11.87
- Very good loaf volume and crumb grain.
- out of mixer=sl. tough; Very low absorption
- Nice Break, Little Dry
- avg. mixt; avg. abs.; nice grain; nice dough; avg. volume; crumb grain=closed

# COOPERATOR'S COMMENTS (con't.)

#### (Small Scale) Kansas

#### 98-410 KS84063-2W

- excellent but very strong dough
- color=white
- Extremely strong flour, open, irregular grain, above average volume
- Excellent exterior and interior crumb, white, silky
- 38 ppm AA; crumb grain
- MOISTURE 13.60 ASH(14%) 0.460 PROTEIN(14%) 14.00
- out of mixer= sl. tough; Poor texture and color
- Nice Break, Large Volume, No Shell Top at all
- sl. long mixt; hi abs.; avg. grain; silky texture; white; nice dough; hi volume.

#### 98-411 KS85W663-42W

- excellent but very strong dough
- crumb grain=sl. dense;
- strong flour, open grain, average volumes
- Excellent exterior and interior crumb, white, silky
- 25 ppm AA
- Strong and extensible dough out of the mixer
- MOISTURE 13.76 ASH(14%) 0.436 PROTEIN(14%) 12.59
- at make-up= sl. tough; cell thickness=thin=desirable
- Nice Break
- long mixt; avg. abs.; good grain; elastic dough at makeup; good volume; crumb texture=good
- Short proof time.

U.S. DEPARTMENT OF AGRICULTURE

The following statements are made in accordance with the Privacy Act of

AGRICULTURAL MARKETING SERVICE		1974 (5 U.S.C. 552a) and the Paperwo	rk Reduction Act (PRA) of 1995.			
EXHIBIT E STATEMENT OF THE BASIS OF OWNERS	HIP	Application is required in order to de certificate is to be issued (7 U.S.C. 2 until certificate is issued (7 U.S.C. 2426				
1. NAME OF APPLICANT(S)		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME			
Kansas Agricultural Experiment Station		КЅ95Н167-3	Stanton			
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Coun	try)	5. TELEPHONE (include area code)	6. FAX (include area code)			
Waters Hall		785-532-6147	785-532-6563			
Kansas State University Manhattan, KS 66506  7. PVPO NUMBER  20020050						
8. Does the applicant own all rights to the variety? Mark an "X" in a	ppropr	iate block. If no, please explain.	X YES NO			
9. Is the applicant (individual or company) a U.S. national or U.S. ba	sed co	ompany?	X YES NO			
If no, give name of country		L-				
0. Is the applicant the original owner?	<u>i</u>	() If no, please answer one of the fo	ollowing:			
a. If original rights to variety were owned by individual(s), is (are)	the or	iginal owner(s) a U.S. national(s)?				
YES		O If no, give name of country				
b. If original rights to variety were owned by a company(ies), is(ar	re) the	original owner(s) a U.S. based company	?			
YES	□ N	O If no, give name of country				
1. Additional explanation on ownership (if needed, use reverse for e	xtra sp	pace):				
LEASE NOTE:						
lant variety protection can be afforded only to owners (not licensees) who r	meet or	ne of the following criteria:				
If the rights to the variety are owned by the original breeder, that person rewhich affords similar protection to nationals of the U.S. for the same general			er country, or national of a country			
If the rights to the variety are owned by the company which employed the member country, or owned by nationals of a country which affords similar						
If the applicant is an owner who is not the original owner, both the original	al own	er and the applicant must meet one of the ab	ove criteria.			
he original breeder/owner may be the individual or company who directed	final br	eeding. See Section 41(a)(2) of the Plant V	ariety Protection Act for definition.			
According to the Paperwork Reduction Act of 1995, no persons are required to respond to this information collection is 0581-0055. The time required to compete this information searching existing data sources, gathering and maintaining the data needed, and complete	n collecti	on is estimated to average 10 minutes per response				
The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the (Not all prohibited bases apply to all programs). Persons with disabilities who require alte USDA's TARGET Center at 202-720-2600 (voice and TDD).						

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

STD-470-E (07-97) (Destroy previous editions).
Electronic version designed using WordPerfect InForms by USDA-AMS-IMB.